

UNIVERSAL MAPS AND SURJECTIVE CHARACTERIZATIONS OF COMPLETELY METRIZABLE LC^n -SPACES

A. CHIGOGIDZE AND V. VALOV

(Communicated by James E. West)

ABSTRACT. We construct an n -dimensional completely metrizable $AE(n)$ -space $P(n, \tau)$ of weight $\tau \geq \omega$ with the following property: for any at most n -dimensional completely metrizable space Y of weight $\leq \tau$ the set of closed embeddings $Y \rightarrow P(n, \tau)$ is dense in the space $C(Y, P(n, \tau))$. It is also shown that completely metrizable LC^n -spaces of weight $\tau \geq \omega$ are precisely the n -invertible images of the Hilbert space $\ell_2(\tau)$.

INTRODUCTION

Let \mathcal{Y} be a class of completely metrizable spaces. A space $X \in \mathcal{Y}$ is said to be *strongly \mathcal{Y} -universal* if for any space $Y \in \mathcal{Y}$ the set of closed embeddings $Y \rightarrow X$ is dense in the space $C(Y, X)$ of all continuous maps from Y to X endowed with the limitation topology (a stronger version of this notion under the same name was introduced in [BM]). This property is very important in the theory of manifolds modelled on certain model spaces. Let us recall the corresponding results:

- (i) If \mathcal{R} is the class of all metrizable compacta, then X is homeomorphic to the Hilbert cube Q iff X is a strongly \mathcal{R} -universal AE -compactum $[T_1]$;
- (ii) if \mathcal{R}_n is the class of all at most n -dimensional metrizable compacta, then X is homeomorphic to Menger's universal n -dimensional compactum M_n^{2n+1} [E] iff X is a strongly \mathcal{R}_n -universal $AE(n)$ -compactum (for $n = 0$, [Br]; for $n \geq 1$, [B]);
- (iii) if \mathcal{M}_τ is the class of all completely metrizable spaces of weight $\leq \tau$, $\tau \geq \omega$, then X is homeomorphic to the Hilbert space $\ell_2(\tau)$ iff X is a strongly \mathcal{M}_τ -universal AE -space $[T_2]$.
- (iv) if $\mathcal{M}_{0,\tau}$ is the class of all zero-dimensional completely metrizable spaces of weight $\leq \tau$, $\tau \geq \omega$, then X is homeomorphic to the Baire space $B(\tau)$ iff X is strongly $\mathcal{M}_{0,\tau}$ -universal (for $\tau = \omega$ [AU]; for $\tau > \omega$, [St]).

Received by the editors August 31, 1987 and, in revised form, June 8, 1989.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 54C55, 54E50, 54F35.

Key words and phrases. Strongly (n, τ) -universal map, n -soft map. $AE(n, m)$ -space.

The second author was partially supported by the Bulgarian Ministry of Culture, Science and Education, Contract N 45.

© 1990 American Mathematical Society
0002-9939/90 \$1.00 + \$.25 per page

The main purpose of this paper is to show the existence of a strongly $\mathcal{M}_{n,\tau}$ -universal $AE(n)$ -space, where $\mathcal{M}_{n,\tau}$ is the class of all at most n -dimensional completely metrizable spaces of weight $\leq \tau$, $\tau \geq \omega$. Let us note that strongly $\mathcal{M}_{n,\omega}$ -universal $AE(n)$ -spaces were constructed by the first author in [C₃]. There are many reasons to hope that the following problems have affirmative solutions:

Problem. Are any two strongly $\mathcal{M}_{n,\tau}$ -universal $AE(n)$ -spaces homeomorphic? In particular, is any strongly $\mathcal{M}_{n,\omega}$ -universal $AE(n)$ -space homeomorphic to Nobeling's universal n -dimensional space N_n^{2n+1} ?

The second part of this paper is devoted to surjective characterizations of completely metrizable LC^n -spaces. Similar characterizations in the class of metrizable compacta were earlier obtained by Hoffman [H₂] and Dranishnikov [D] (see also [C₄], where the class of Polish spaces is considered).

1. PRELIMINARIES

All spaces considered are metrizable and maps continuous. By dimension \dim we mean covering dimension. A metrizable space X is an *absolute (neighborhood) extensor* in dimension n (briefly, $X \in A(N)E(n)$) if for any at most n -dimensional metrizable space Y and any closed subspace A of it each map $f: A \rightarrow X$ can be extended to the whole of Y (respectively, to a neighborhood of A in Y). It is well known that for $n > 0$, $X \in A(N)E(n)$ iff $X \in LC^{n-1} \cap C^{n-1}$ (respectively, $X \in LC^{n-1}$). Note also that any metrizable space, and hence any completely metrizable space, is an $AE(0)$. (The argument can be made as follows: Let $\dim Y = 0$ and $A \subset Y$ be closed. Let X be metrizable and $f: A \rightarrow X$ be a map. It is well known that there is a retraction $r: Y \rightarrow A$. Then $f \circ r$ is the desired extension of f .) The notion of *n -soft map* between compacta was introduced by Schepin [S]. Later Chigogidze [C₁] extended it to the class of all Tychonov spaces. Below we use the following definition of this notion: a map $f: X \rightarrow Y$ between metrizable spaces is called *n -soft* if for any at most n -dimensional paracompact space Z , any closed subspace A of it and any two maps $g: Z \rightarrow Y$, $h: A \rightarrow X$ with $f \circ h = g|_A$, there exists a map $k: Z \rightarrow X$ such that $f \circ k = g$ and $k|_A = h$.

2. UNIVERSAL MAPS

Lemma 2.1. *Let $f: X \rightarrow Y$ be an n -soft map between metrizable spaces. Suppose $\dim X \leq n$ and Y is a complete absolute extensor for the class of all metrizable spaces. Then X is complete.*

Proof. Consider the Stone-Ćech compactification βX of X . Denote by Z the space obtained from βX by means of making the points of $\beta X - X$ isolated. This space is shown to be paracompact in the proof of [P, Lemma 2].

Claim. $\dim Z \leq n$. Since Z is normal it suffices to extend to Z an arbitrary map $g: F \rightarrow S^n$ from a closed subset F of Z into the n -dimensional sphere.

The case $F \subseteq Z - X$ is trivial. Suppose now that $F \cap X \neq \emptyset$. Since $\dim X \leq n$, there exists an extension $g_1: F \cup X \rightarrow S^n$ of g . Observe that $F \cup X$ is closed in Z . Hence we can extend g_1 to a map g_2 from Z into the $(n+1)$ -dimensional disk B^{n+1} . Put $H = g_2^{-1}(B^{n+1} - \{b\})$, where $b \in B^{n+1} - S^n$. Fix a retraction $r: (B^{n+1} - \{b\}) \rightarrow S^n$. Clearly, H is clopen in Z ; so there exists a map $g_3: Z \rightarrow S^n$ extending the composition $r \circ g_2: H \rightarrow S^n$. Obviously, $g_3|_F = g$. The claim is proved.

Since Y is metrizable and is an AE for metrizable spaces, then Y is an absolute retract for metrizable spaces. Further, complete metrizable implies Čech complete. Being metrizable, Y is a paracompact p -space, so using [P, Fact 6 and Corollary 1(b)], we conclude that Y is an absolute extensor for the class of collectionwise normal spaces. Take a map $h: Z \rightarrow Y$ such that $h|_X = f$. It follows by the n -softness of f that there exists a retraction from Z onto X . Now, by arguments of Przymusiński ([P, the proof of Lemma 2]), X is complete.

Lemma 2.2 (for $\tau = \omega$, $[C_2]$). *Let $0 \leq n < \omega \leq \tau$. Then there exist an n -dimensional completely metrizable space X of weight τ and an n -soft map $f: X \rightarrow \ell_2(\tau)$.*

Proof. By $[C_2$, Theorem 5], we can suppose that $\tau > \omega$. Let d_1 be any metric on $\ell_2(\tau)$. Fix a completely zero-dimensional (with respect to d_1) surjection $g: \ell_2(\tau) \rightarrow Y$, where Y is a separable metrizable space [AP]. By an application of $[C_2$, Theorem 5.1] (see also the remarks following it) there exist an at most n -dimensional separable space Z and an n -soft map $h: Z \rightarrow Y$. Let X be a fibered product (pullback) of the spaces $\ell_2(\tau)$ and Z with respect to the maps g and h . Denote by $f: X \rightarrow \ell_2(\tau)$ and $p: X \rightarrow Z$ the corresponding canonical projections. Let d_2 be any metric on Z . Clearly, the map p is completely zero-dimensional with respect to the metric $d = (d_1^2 + d_2^2)^{1/2}$ [AP, Chapter 6, §3, Lemma 4]. Hence $\dim X \leq \dim Z \leq n$. It is easy to see that $\omega(X) = \tau$. Observe also that the n -softness of h implies the n -softness of f . Since $\ell_2(\tau)$ contains a copy of the n -dimensional cube I^n and f is an n -soft map, the space X contains a copy of I^n too. Thus $\dim X = n$. By Lemma 2.1, X is completely metrizable.

Corollary 2.3 (for $\tau = \omega$, $[C_3]$). *Let $0 \leq n < \omega \leq \tau$. Then for every (completely) metrizable space Y of weight τ there exist an at most n -dimensional (completely) metrizable space Z of weight τ and an n -soft map $g: Z \rightarrow Y$.*

Proof. Embed Y into $\ell_2(\tau)$ as a (closed) subspace and consider the map $f: X \rightarrow \ell_2(\tau)$ from Lemma 2.2. Put $Z = f^{-1}(Y)$ and $g = f|_Z$.

Definition 2.4 (for $\tau = \omega$, $[C_3]$). Let $0 \leq n < \omega \leq \tau$. A map $f: X \rightarrow Y$ is said to be (n, τ) -full if for any map $g: Z \rightarrow Y$ from any at most n -dimensional completely metrizable space Z of weight $\leq \tau$ there exists a closed embedding $h: Z \rightarrow X$ such that $f \circ h = g$.

Definition 2.5 (for $\tau = \omega, [C_3]$). Let $0 \leq n < \omega \leq \tau$. A map $f: X \rightarrow Y$ is called strongly (n, τ) -universal if for any open cover \mathcal{U} of X , any at most n -dimensional completely metrizable space Z of weight $\leq \tau$ and any map $g: Z \rightarrow X$ there exists a closed embedding $h: Z \rightarrow X$ \mathcal{U} -close to g with $f \circ h = f \circ g$. We shall also say that a space X is strongly (n, τ) -universal if the constant map $X \rightarrow *$ is strongly (n, τ) -universal in the above sense.

Lemma 2.6 (for $\tau = \omega, [C_3]$). Let $0 \leq n < \omega \leq \tau$ and $S = \{X_k, p_k^{k+1}, \omega\}$ be an inverse sequence consisting of completely metrizable spaces X_k of weight $\leq \tau$ and n -soft, (n, τ) -full projections p_k^{k+1} . Then the limit projection $p_0: X \rightarrow X_0$, where $X = \lim S$, is strongly (n, τ) -universal.

Proof. Equip X with the metric $d(\{x_k\}, \{y_k\}) = \max_k d_k(x_k, y_k)$, where d_k is a metric for X_k with $d_k \leq 2^{-k}$, $k \in \omega$. It suffices to show that, given a completely metrizable space Y with $\dim Y \leq n$ and $\omega(Y) \leq \tau$ and maps $f: Y \rightarrow X$, $\alpha: X \rightarrow (0, 1)$, there is a closed embedding $g: Y \rightarrow X$ with $p_0 \circ g = p_0 \circ f$ and $d(f(y), g(y)) \leq \alpha(f(y))$ for each $y \in Y$.

For each $k \in \omega$ fix a closed embedding $i_{k+1}: X_{k+1} \rightarrow \ell_2(\tau)$. Since $\ell_2(\tau) \in AE$ there is a map $h: (\ell_2(\tau))^2 \times [0, \infty) \rightarrow \ell_2(\tau)$ such that $h(a, b, t) = a$ for $t \leq 1$ and $h(a, b, t) = b$ for $t \geq 2$. By Corollary 2.3, for each $k \in \omega$ there is an n -soft map $q_{k+1}: Z_{k+1} \rightarrow X_k \times \ell_2(\tau)$, where Z_{k+1} is a completely metrizable space with $\dim Z_{k+1} \leq n$ and $\omega(Z_{k+1}) = \tau$. The n -softness of the projection p_k^{k+1} implies the existence of a $r_{k+1}: Z_{k+1} \rightarrow X_{k+1}$ with $p_k^{k+1} \circ r_{k+1} = \pi_k \circ q_{k+1}$ and $r_{k+1}|_{A_{k+1}} = (p_k^{k+1} \Delta i_{k+1})^{-1} \circ q_{k+1}|_{A_{k+1}}$, where

$$A_{k+1} = q_{k+1}^{-1}((p_k^{k+1} \Delta i_{k+1})(X_{k+1})) \quad \text{and} \quad \pi_k: X_k \times \ell_2(\tau) \rightarrow X_k$$

denotes the natural projection. Put $g_0 = p_0 \circ f$. By our assumption, the projection p_0^1 of the spectrum S is (n, τ) -full. Hence there exists a closed embedding $j_1: Y \rightarrow X_1$ such that $p_0^1 \circ j_1 = g_0$. Consider now the map

$$g_0 \Delta h(i_1 \circ p_1 \circ f \Delta i_1 \circ j_1 \Delta 2\alpha \circ f): Y \rightarrow X_0 \times \ell_2(\tau).$$

Since $\dim Y \leq n$ and the map q_1 is n -soft there is a map $s_1: Y \rightarrow Z_1$ such that $g_0 \Delta h(i_1 \circ p_1 \circ f \Delta i_1 \circ j_1 \Delta 2\alpha \circ f) = q_1 \circ s_1$. We define a map $g_1: Y \rightarrow X_1$ by the formula $g_1 = r_1 \circ s_1$. Note that $p_0^1 \circ g_1 = g_0$. If $y \in Y$ and $\alpha(f(y)) \leq 2^{-1}$, then $g_1(y) = p_1(f(y))$. (Use the fact that $g_1(y) = r_1(s_1(y))$, show that $s_1(y) \in A_1$, and then use the above formula for $q_1 \circ s_1$.)

Let us construct a map $g_2: Y \rightarrow X_2$. By our assumption, the projection p_1^2 is (n, τ) -full. Hence there is a closed embedding $j_2: Y \rightarrow X_2$ such that $p_1^2 \circ j_2 = g_1$. Since $\dim Y \leq n$ and the map q_2 is n -soft there is a map $s_2: Y \rightarrow Z_2$ such that $q_2 \circ s_2 = g_1 \Delta h(i_2 \circ p_2 \circ f \Delta i_2 \circ j_2 \Delta 2^2 \alpha \circ f)$. Put $g_2 = r_2 \circ s_2$. As above, $p_1^2 \circ g_2 = g_1$. Note that if $y \in Y$ and $\alpha(f(y)) \leq 2^{-2}$, then $g_2(y) = p_2(f(y))$; if $\alpha(f(y)) \geq 2^{-1}$, then $g_2(y) = j_2(y)$.

Let us suppose that for each i , $2 \leq i \leq k$, we have already constructed maps $g_i: Y \rightarrow X_i$ and closed embeddings $j_i: Y \rightarrow X_i$ satisfying the following conditions:

- (1) _{i} $p_{i-1}^i \circ g_i = g_{i-1}$;
- (2) _{i} $p_{i-1}^i \circ j_i = g_{i-1}$;
- (3) _{i} if $y \in Y$ and $\alpha(f(y)) \leq 2^{-i}$, then $g_i(y) = p_i(f(y))$;
- (4) _{i} if $y \in Y$ and $\alpha(f(y)) \leq 2^{-i+1}$, then $g_i(y) = j_i(y)$.

Let us construct a map $g_{k+1}: Y \rightarrow X_{k+1}$ and a closed embedding $j_{k+1}: Y \rightarrow X_{k+1}$ with the desired properties. Choose an arbitrary closed embedding $j_{k+1}: Y \rightarrow X_{k+1}$ with $p_k^{k+1} \circ j_{k+1} = g_k$ (the existence of j_{k+1} follows from the (n, τ) -fullness of p_k^{k+1}). As above there is a map $s_{k+1}: Y \rightarrow Z_{k+1}$ such that $q_{k+1} \circ s_{k+1} = g_k \Delta h(i_{k+1} \circ p_{k+1} \circ f \Delta i_{k+1} \circ j_{k+1} \Delta 2^{k+1} \alpha \circ f)$. We put $g_{k+1} = r_{k+1} \circ s_{k+1}$. The verification of the conditions (1) _{$k+1$} —(4) _{$k+1$} is left to the reader.

It follows from the conditions (1) _{k} , $k \in \omega$, that the diagonal product $g = \Delta\{g_k: k \in \omega\}$ maps Y into X and satisfies the following equalities: $p_k \circ g = g_k$, $k \in \omega$. In particular, we have $p_0 \circ g = g_0 = p_0 \circ f$.

Let $y \in Y$. If $\alpha(f(y)) \in [2^{-k-1}, 2^{-k}]$, then $2^i \alpha(f(y)) \leq 1$ for $i \leq k$; hence

$$p_i(f(y)) = g_i(y) \quad \text{for } i \leq k$$

and

$$\begin{aligned} d(f(y)g(y)) &= \max\{d_i(p_i(f(y)), g_i(y)): i = k + 1, k + 2, \dots\} \\ &\leq 2^{-k-1} \leq \alpha(f(y)). \end{aligned}$$

In order to show that g is a closed embedding it suffices to use the corresponding arguments from the proof of [C₂, Lemma 7.11].

Theorem 2.7 (for $\tau = \omega$, [C₃]). *Let $0 \leq n < \omega \leq \tau$. Then there exist an n -dimensional completely metrizable space $P(n, \tau)$ of weight τ and a strongly (n, τ) -universal n -soft map $f(n, \tau): P(n, \tau) \rightarrow \ell_2(\tau)$.*

Proof. Put $X_0 = \ell_2(\tau)$. By Corollary 2.3, there exists at most an n -dimensional completely metrizable space X_{k+1} of weight τ and an n -soft map $h_{k+1}: X_{k+1} \rightarrow X_k \times \ell_2(\tau)$. Put $p_k^{k+1} = \pi_k \circ h_{k+1}$, where $\pi_k: X_k \times \ell_2(\tau) \rightarrow X_k$ is the natural projection, $k \in \omega$. So we get an inverse sequence $S = \{X_k, p_k^{k+1}\}$ consisting of at most n -dimensional completely metrizable spaces of weight τ and (n, τ) -full n -soft projections. Put $P(n, \tau) = \lim S$ and $f(n, \tau) = p_0$. By Lemma 2.6, the map $f(n, \tau)$ is strongly (n, τ) -universal. Obviously, this map is n -soft. Since $\dim X_k \leq n$ for each $k \geq 1$, we have $\dim P(n, \tau) \leq n$ [N]. The inverse inequality $\dim P(n, \tau) \geq n$ follows from the strong (n, τ) -universality of $f(n, \tau)$. Finally observe that $P(n, \tau)$ is a complete metrizable space of weight τ . This completes the proof.

Corollary 2.8 (for $\tau = \omega$, $[C_3]$). *Let $0 \leq n < \omega \leq \tau$. The space $P(n, \tau)$ is an n -dimensional strongly (n, τ) -universal completely metrizable $AE(n)$ -space of weight τ .*

Proof. $P(n, \tau)$ is an $AE(n)$ -space as an n -soft preimage of $\ell_2(\tau)$.

Remark 2.9. The space $P(n, \tau)$ has the following property: for every open subspace U of $P(n, \tau)$ and any at most n -dimensional completely metrizable space X of weight $\leq \tau$ there exists an embedding $h: X \rightarrow U$ such that $h(X)$ is closed in $P(n, \tau)$. Indeed, consider a constant map $g: X \rightarrow \{p\}$, where p is an arbitrary point of U , and the open cover $\mathcal{U} = \{U, P(n, \tau) - \{p\}\}$ of $P(n, \tau)$. By Definition 2.5, there is a closed embedding $h: X \rightarrow P(n, \tau)$ \mathcal{U} -close to g . Clearly, $h(X) \subset U$. In particular, we have $\omega(U) = \tau$ for every open subset U of $P(n, \tau)$. Hence, $P(0, \tau)$ is homeomorphic to the Baire space $B(\tau)$ (see [St]).

Remark 2.10. A. Wasko proved, in [W], that for every $n \geq 0$ and every $\tau \geq \omega$ there exists an n -dimensional completely metrizable space $X_{n, \tau}$ of weight τ such that every at most n -dimensional completely metrizable space of weight $\leq \tau$ is embedded in $X_{n, \tau}$ as a closed subset (for $\tau = \omega$ this was proved earlier by the first author, see $[C_1]$, Corollary 3). Recently E. Pol, [Po], strengthened this result of A. Wasko by proving that for every n -dimensional completely metrizable space X of weight $\leq \tau$ the set of all embeddings of X onto a closed subset of $S(\tau)^\omega$ contained in $K_n(\tau)$ is residual in the space $C(X, S(\tau)^\omega)$. Here $S(\tau)$ is the τ -star-space and $K_n(\tau)$ denotes Nagata's universal n -dimensional space.

3. SURJECTIVE CHARACTERIZATIONS OF LC^n -SPACES

Definition 3.1 $[H_2]$. A space X is said to be in the class $AE(n, m)$, where $0 \leq n \leq m \leq \infty$, if for any metrizable space Z with $\dim Z \leq m$, any closed subspace A of it with $\dim A \leq n$, any map $f: A \rightarrow X$ can be extended to the whole of Z . If f can be extended only to a neighborhood of A in Z , we get a definition of the class $ANE(n, m)$.

Lemma 3.2. *For every $n \in \omega$ the following equalities are true in the class of all metrizable spaces: $A(N)E(n+1) = A(N)E(n, n+1)$.*

In the class of all metrizable compacta this lemma was proved by Dranishnikov (see [D, Lemma 3.1]). The same proof remains valid in the general case.

Lemma 3.3. *If $0 \leq n < m \leq \infty$, then the following equalities are true in the class of all metrizable spaces: $A(N)E(n+1) = A(N)E(n, m)$.*

Proof. Since $n+1 \leq m$, then $A(N)E(n, m) \subset A(N)E(n, n+1)$; thus, $A(N)E(n, m) \subset A(N)E(n+1)$ follows from Lemma 3.2. Suppose X is a metrizable $AE(n+1)$ -space. Let Z be a metrizable space with $\dim Z \leq m$, A a closed subspace of Z with $\dim A \leq n$ and f a map from A to X . Take a metrizable AE -space, say Y , of dimension $\leq n+1$ containing A as a closed

subspace ([K], see also [C₄] for Polish spaces and [Ts] for completely metrizable spaces). Now choose a map $k: Y \rightarrow X$ such that $k|_A = f$. This is possible because, by Lemma 3.2, $X \in AE(n, n + 1)$. Since $Y \in AE$, there exists a map $g: Z \rightarrow Y$ such that $g|_A = id$. Then $k \circ g$ is an extension of f . The inclusion $ANE(n + 1) \subset ANE(n, m)$ follows from the same arguments.

Definition 3.4 [H₁]. A map $f: X \rightarrow Y$ is called n -invertible if for any at most n -dimensional metrizable space Z and any map $g: Z \rightarrow Y$ there exists a map $h: Z \rightarrow X$ such that $g = f \circ h$.

Obviously, every n -invertible map is a surjection. For a metrizable compactum X it is known [D], [H₂] that X is an $AE(n + 1)$ iff X is an n -invertible image of Q . It is also true [D] that the class of all metrizable $AE(n)$ -compacta coincides with the class of n -invertible images of the n -dimensional universal Menger compactum m_n^{2n+1} . The first author [C₄] gave similar characterizations of $AE(n + 1)$ and $AE(n)$ in the class of Polish spaces. In the case of metrizable spaces of uncountable weight we know only the following facts: (i) the classical characterization of completely metrizable spaces of weight τ as open images of the Baire space $B(\tau)$; (ii) a metrizable space X of weight τ is Čech-complete iff X is a 0-invertible image of $B(\tau)$ [V].

Below we give similar characterizations of metrizable $AE(n)$ -spaces of arbitrary weight.

Definition 3.5 [C₄]. A map $f: X \rightarrow Y$ is said to be inductively n -soft if there exists a closed subspace Z of X such that the restriction $f|_Z: Z \rightarrow Y$ is n -soft.

Theorem 3.6. *Let X be a metrizable space of weight $\tau \geq \omega$. Then for every $n \in \omega$ the following conditions are equivalent:*

- (i) $X \in AE(n + 1)$ (respectively, $X \in ANE(n + 1)$);
- (ii) X is an inductively n -soft image of an AE (respectively, of an ANE);
- (iii) X is an n -invertible image of an AE (respectively, of an ANE);

Proof. We shall prove only the global variant. The local one follows from the same arguments.

(i) \rightarrow (ii). By Corollary 2.3, there exist an at most n -dimensional metrizable space Y of weight τ and an n -soft map $g: Y \rightarrow X$. Embed Y into a metrizable AE -space Z as a closed subspace. By Lemma 3.3, there exists an extension $h: Z \rightarrow X$ of g . Clearly, h is inductively n -soft.

(ii) \rightarrow (iii). This implication is trivial, because any n -soft map is n -invertible.

(iii) \rightarrow (i). Let Z be an AE -space and $f: Z \rightarrow X$ be an n -invertible map. In view of Lemma 3.2 it suffices to show that $X \in AE(n, n + 1)$. Let B be any at most $(n + 1)$ -dimensional metrizable space, A a closed subspace of B with $\dim A \leq n$ and g a map from A to X . Since f is n -invertible, there exists a map $h: A \rightarrow Z$ such that $f \circ h = g$. Take any extension $k: B \rightarrow Z$ of h . Then the map $f \circ k$ is an extension of g .

Let us consider the proof of Theorem 3.6. If X is a complete metrizable space of weight τ , then the space Y is also complete, so we can suppose that Z is the space $\ell_2(\tau)$. Thus, the following theorem is true.

Theorem 3.7. *Let X be a completely metrizable space of weight $\tau \geq \omega$. Then for every $n \in \omega$ the following conditions are equivalent:*

- (i) $X \in AE(n+1)$ (respectively, $X \in ANE(n+1)$);
- (ii) X is an inductively n -soft image of $\ell_2(\tau)$ (respectively, of an open subspace of $\ell_2(\tau)$);
- (iii) X is an n -invertible image of $\ell_2(\tau)$ (respectively, of an open subset of $\ell_2(\tau)$).

The proof of the following result is analogous to the proof of Theorem 3.6.

Theorem 3.8. *Let X be a completely metrizable space of weight $\tau \geq \omega$. Then for every $n \in \omega$ the following conditions are equivalent:*

- (i) $X \in AE(n)$ (respectively, $X \in ANE(n)$);
- (ii) X is an inductively n -soft image of $P(n, \tau)$ (respectively, of an open subset of $P(n, \tau)$);
- (iii) X is an n -invertible image of $P(n, \tau)$ (respectively, of an open subset of $P(n, \tau)$).

ACKNOWLEDGMENTS

The authors express their sincere appreciation to the referee and to W. Olszewski for many helpful suggestions.

REFERENCES

- [AP] P. Alexandrov and B. Pasynkov, *Introduction to dimension theory*, Nauka, Moscow 1973. (Russian)
- [AU] P. Alexandrov and P. Urysohn, *Über nulldimensionale mengen*, Math. Ann. **98** (1928), 89–106.
- [B] M. Bestvina, *Characterizing k -dimensional universal Menger compacta*, Dissertation, Knoxville, 1984.
- [BM] M. Bestvina and J. Mogilski, *Characterizing certain incomplete infinite-dimensional absolute retracts*, Michigan Math. J. **33** (1986), 291–313.
- [Br] L. Brouwer, *On the structure of perfect sets of points*, Proc. Akad. Amsterdam **12** (1910), 785–794.
- [C₁] A. Chigogidze, *Uncountable powers of the real line and the natural series and n -soft maps*, Dokl. Akad. Nauk SSSR **278** (1984), 50–53. (Russian)
- [C₂] —, *Noncompact absolute extensors in dimension n , n -soft maps and their applications*, Izvestia Akad. Nauk SSSR, Ser. Math. **50** (1986), 156–180. (Russian)
- [C₃] —, *n -soft maps of n -dimensional spaces*, Mat. Zametki, **46** (1989), 88–95. (Russian)
- [C₄] —, *Characterization of Polish $AE(n)$ -spaces*, Vestnik Mosk. Gos. Univ. **5** (1987), 32–35. (Russian)
- [D] A. Dranishnikov, *Absolute extensors in dimension n and n -soft maps increasing dimension*, Uspekhi Mat. Nauk **39** (1984), 55–95. (Russian)

- [E] R. Engelking, *Dimension theory*, PWN, Warsaw 1978.
- [H₁] B. Hoffman, *A surjective characterization of Dugundji spaces*, Proc. Amer. Math. Soc. **76** (1979), 151–156.
- [H₂] —, *An injective characterization of Peano spaces*, Top. Appl. **11** (1980), 37–46.
- [K] Y. Kodama, *On embeddings of spaces into ANR and shape*, J. Math. Soc. Japan **27** (1975), 533–544.
- [N] K. Nagami, *Finite-to-one closed mappings and dimension II*, Proc. Japan Acad. **35** (1959), 437–439.
- [P] T. Przymusiński, *Collectionwise normality and absolute retracts*, Fund. Math. **98** (1978), 61–73.
- [Po] E. Pol, *Residuality of the set of embeddings into Nagata's n -dimensional universal space*, Fund. Math. **129** (1988), 59–67.
- [S] E. Shchepin, *Functors and uncountable powers of compacta*, Uspekhi Mat. Nauk **36** (1981), 3–62. (Russian)
- [St] A. Stone, *Non-separable Borel sets*, Rozprawy Mat. **28** (1962), 1–40.
- [T₁] H. Toruńczyk, *On CE -images of the Hilbert cube and characterization of Q -manifolds*, Fund. Math. **106** (1980), 31–40.
- [T₂] —, *Characterizing Hilbert space topology*, Fund. Math. **111** (1981), 247–262.
- [Ts] K. Tsuda, *A note on closed embeddings of finite dimensional metric spaces*, Bull. Polish Acad. Sci. Math. **33** (1985), 541–546.
- [V] V. Valov, *Milutin mappings and $AE(0)$ -spaces*, Comp. Rend. Bulg. Acad. Sci. **40** (1987), 9–12.
- [W] A. Wasko, *Spaces universal under closed embeddings for finite dimensional complete metric spaces*, Bull. London Math. Soc. **18** (1986), 293–298.

DEPARTMENT OF MATHEMATICS, MOSKOW STATE UNIVERSITY, MOSKOW 119899, USSR

DEPARTMENT OF MATHEMATICS, SOFIA STATE UNIVERSITY, SOFIA 1126, BULGARIA